

List of publications

Taina Pihlajaniemi

1 April, 2016

A Peer-reviewed scientific articles

Journal article (refereed), original research; review article, literature review, systematic review; book section, chapters in research books; conference proceedings

- A.1 Savolainen E-R, Kero M, Pihlajaniemi T, and Kivirikko KI. Deficiency of galactosylhydroxyllysyl glucosyltransferase, an enzyme of collagen synthesis, in a family with dominant epidermolysis bullosa simplex. *N Engl J Med* 304: 197-204, 1981
- A.2 Pihlajaniemi T, Myllylä R, Alitalo K, Vaheri A, and Kivirikko KI. Post-translational modifications in the biosynthesis of type IV collagen by a human tumor cell line. *Biochemistry* 20: 7409-7415, 1981
- A.3 Anttinen H, Puistola U, Pihlajaniemi T, and Kivirikko KI. Differences between proline and lysine hydroxylations in their inhibition by zinc or by ascorbate deficiency during collagen synthesis in various cell types. *Biochim Biophys Acta* 674: 336-344, 1981
- A.4 Pihlajaniemi T, Myllylä R, Kivirikko KI, and Tryggvason K. Effects of streptozotocin diabetes, glucose, and insulin on the metabolism of type IV collagen and proteoglycan in murine basement membrane-forming EHS tumor tissue. *J Biol Chem* 257: 14914-14920, 1982
- A.5 de Wet WJ, Pihlajaniemi T, Myers J, Kelly TE, and Prockop DJ. Synthesis of a shortened pro- α 2(I) chain and decreased synthesis of pro- α 2(I) chains in a proband with osteogenesis imperfecta. *J Biol Chem* 258: 7721-7728, 1983
- A.6 Oikarinen J, Pihlajaniemi T, Hämäläinen L, and Kivirikko KI. Cortisol decreases the cellular concentration of translatable procollagen mRNA species in cultured human skin fibroblasts. *Biochim Biophys Acta* 741: 297-302, 1983
- A.7 Myllylä R, Koivu J, Pihlajaniemi T, and Kivirikko KI. Protein disulphide-isomerase activity in various cells synthesizing collagen. *Eur J Biochem* 134: 7-11, 1983
- A.8 Dickson LA, Pihlajaniemi T, Deak S, Pope FM, Nicholls A, Prockop DJ, and Myers JC. Nuclease S1 mapping of a homozygous mutation in the carboxyl-propeptide-coding region of the pro α 2(I) collagen gene in a patient with osteogenesis imperfecta. *Proc Natl Acad Sci USA* 81: 4524-4528, 1984
- A.9 Pihlajaniemi T, Dickson LA, Pope FM, Korhonen VR, Nicholls A, Prockop DJ, and Myers JC. Osteogenesis imperfecta: cloning of a pro- α 2(I) collagen gene with a frameshift mutation. *J Biol Chem* 259: 12941- 12944, 1984
- A.10 Loidl HR, Brinker JM, May M, Pihlajaniemi T, Morrow S, Rosenbloom J, and Myers JC. Molecular cloning and carboxyl-propeptide analysis of human type III procollagen. *Nucleic Acids Res* 12: 9383-9394, 1984

- A.11 Pihlajaniemi T, Tryggvason K, Myers JC, Kurkinen K, Lebo R, Cheung M- C, Prockop DJ, and Boyd CD. cDNA clones coding for the pro- α 1(IV) chain of human type IV procollagen reveal an unusual homology of amino acid sequences in two halves of the carboxyl-terminal domain. *J Biol Chem* 260: 7681-7687, 1985
- A.12 Myers JC, Dickson LA, Pope FM, Korhonen VR, Nicholls A, Prockop DJ, and Pihlajaniemi T. A homozygous frameshift mutation in the pro- α 2(I) collagen COOH-propeptide results in osteogenesis imperfecta. *Ann NY Acad Sci* 460: 482-485, 1985
- A.13 Prockop DJ, Chu M-L, de Wet W, Myers JC, Pihlajaniemi T, Ramirez F, and Sippola M. Mutations in osteogenesis imperfecta leading to the synthesis of abnormal type I procollagen. *Ann NY Acad Sci* 460: 289- 297, 1985
- A.14 Soininen R, Tikka L, Chow L, Pihlajaniemi T, Kurkinen M, Prockop DJ, Boyd DC, and Tryggvason K. Large introns in the 3' end of the gene for the pro α 1(IV) chain of human basement membrane collagen. *Proc Natl Acad Sci USA* 83: 1568-1572, 1986
- A.15 Uitto J, Matsuoka LY, Chu M-L, Pihlajaniemi T, and Prockop DJ. Connective tissue biochemistry of neurofibromas. *Ann NY Acad Sci* 486: 271-286, 1986
- A.16 Pihlajaniemi T, Helaakoski T, Tasanen K, Myllylä R, Huhtala M-L, Koivu J, and Kivirikko KI. Molecular cloning of the β -subunit of human prolyl 4-hydroxylase. This subunit and protein disulphide isomerase are products of the same gene. *EMBO J* 6: 643-649, 1987
- A.17 Pihlajaniemi T, Myllylä R, Seyer J, Kurkinen M, and Prockop DJ. Partial characterization of a low molecular weight human collagen that undergoes alternative splicing. *Proc Natl Acad Sci USA* 84: 940-944, 1987
- A.18 Koivu J, Myllylä R, Helaakoski T, Pihlajaniemi T, Tasanen K, and Kivirikko KI. A single polypeptide acts both as the β subunit of prolyl 4-hydroxylase and as a protein disulfide-isomerase. *J Biol Chem* 262: 6447-6449, 1987
- A.19 Kurkinen M, Condon MR, Blumberg B, Barlow DP, Quinones S, Saus J, and Pihlajaniemi T. Extensive homology between the carboxyl-terminal peptides of mouse α 1(IV) and α 2(IV) collagen. *J Biol Chem* 262: 8496- 8499, 1987
- A.20 Pikkarainen T, Eddy R, Fukushima Y, Byers M, Shows T, Pihlajaniemi T, Saraste M, and Tryggvason K. Human laminin B1 chain. A multidomain protein with gene (LAMB1) locus in the q22 region of chromosome 7. *J Biol Chem* 262: 10454-10462, 1987
- A.21 Ala-Kokko L, Pihlajaniemi T, Myers JC, Kivirikko KI, and Savolainen E-R. Gene expression of type I, III and IV collagens in hepatic fibrosis induced by dimethylnitrosamine in the rat. *Biochem J* 244: 75-79, 1987
- A.22 Pihlajaniemi T, and Myers JC. Characterization of a pro- α 2(I) collagen gene mutation by nuclease S1 mapping. *Methods Enzymol* 145: 213-222, 1987
- A.23 Tikka L, Pihlajaniemi T, Henttu P, Prockop DJ, and Tryggvason K. Unique gene structure for the α 1 chain of a human short-chain collagen (type XIII) with alternatively spliced transcripts and translation termination codon at the 5'end of the last exon. *Proc Natl Acad Sci USA* 85: 7491-7495, 1988

- A.24 Tasanen K, Parkkonen T, Chow LT, Kivirikko KI, and Pihlajaniemi T. Characterization of the human gene for a polypeptide that acts both as the β subunit of prolyl 4-hydroxylase and as protein disulfide isomerase. *J Biol Chem* 263: 16218-16224, 1988
- A.25 Pajunen L, Myllylä R, Helaakoski T, Pihlajaniemi T, Tasanen K, Höyhty M, Tryggvason K, Solomon E, and Kivirikko KI. Assignment of the gene coding for both the β -subunit of prolyl 4-hydroxylase and the enzyme protein disulfide isomerase to human chromosome region 17p11qter. *Cytogenet Cell Genet* 47: 37-41, 1988
- A.26 Parkkonen T, Kivirikko KI, and Pihlajaniemi T. Molecular cloning of a multifunctional chicken protein acting as the prolyl 4-hydroxylase β -subunit, protein disulfide-isomerase and a cellular thyroid-hormone-binding protein. Comparison of cDNA-deduced amino acid sequences with those in other species. *Biochem J* 256: 1005-1011, 1988
- A.27 Sandberg M, Tamminen M, Hirvonen H, Vuorio E, and Pihlajaniemi T. Expression of mRNAs coding for the $\alpha 1$ chain of type XIII collagen in human fetal tissues: comparison with expression of mRNAs for collagen types I, II and III. *J Cell Biol* 109: 1371-1379, 1989
- A.28 Helaakoski T, Vuori K, Myllylä R, Kivirikko KI, and Pihlajaniemi T. Molecular cloning of the α -subunit of human prolyl 4-hydroxylase: the complete cDNA-derived amino acid sequence and evidence for alternative splicing of RNA transcripts. *Proc Natl Acad Sci USA* 86: 4392-4396, 1989
- A.29 Saus J, Quinones S, McKrell A, Blumberg B, Muthukumaran G, Pihlajaniemi T, and Kurkinen M. The complete primary structure of mouse $\alpha 2(\text{IV})$ collagen: alignment with mouse $\alpha 1(\text{IV})$ collagen. *J Biol Chem* 264: 6318-6324, 1989
- A.30 Pajunen L, Jones TA, Helaakoski T, Pihlajaniemi T, Sheer D, Solomon E, and Kivirikko KI. Assignment of the gene coding for the α -subunit of prolyl 4-hydroxylase to human chromosome region 10q21.3-23.1. *Am J Hum Genet* 45: 829-834, 1989
- A.31 Pihlajaniemi T, McKeon J, Gay S, Gay R, de Wet WJ, Myers JC, and Prockop DJ. Increased expression of the gene for the pro $\alpha 1(\text{IV})$ chain of basement-membrane procollagen in cultured skin fibroblasts from two variants of osteogenesis imperfecta. *Biochem J* 257: 439-445, 1989
- A.32 Pajunen L, Tamminen M, Solomon E, and Pihlajaniemi T. Assignment of the gene coding for the $\alpha 1$ chain of collagen type XIII to chromosome region 10q11 \rightarrow qter. *Cytogenet Cell Genet* 52: 190-193, 1989
- A.33 Kivirikko KI, Myllylä R, and Pihlajaniemi T. Protein hydroxylation: prolyl 4-hydroxylase, an enzyme with four cosubstrates and a multifunctional subunit. *FASEB J* 3: 1609-1617, 1989
- A.34 Karvonen K, Ala-Kokko L, Pihlajaniemi T, Helaakoski T, Henke S, Günzler V, Kivirikko KI, and Savolainen E-R. Specific inactivation of prolyl 4-hydroxylase and inhibition of collagen synthesis by oxaproline-containing peptides in cultured human skin fibroblasts. *J Biol Chem* 265: 8415-8419, 1990
- A.35 Helaakoski T, Pajunen L, Kivirikko KI, and Pihlajaniemi T. Increases in mRNA concentrations of the α and β -subunits of prolyl 4-hydroxylase accompany increased

gene expression of type IV collagen during differentiation of mouse F9 cells. *J Biol Chem* 265: 11413-11416, 1990

- A.36 Pihlajaniemi T, Pohjolainen E-R, and Myers JC. Complete primary structure of the triple-helical region and carboxyl-terminal domain of a new type IV collagen chain, $\alpha 5(\text{IV})$. *J Biol Chem* 265: 13758-13766, 1990
- A.37 Pihlajaniemi T, and Tamminen M. The $\alpha 1$ chain of type XIII collagen consists of three collagenous and four non-collagenous domains and its primary transcript undergoes complex alternative splicing. *J Biol Chem* 265: 16922-16928, 1990
- A.38 Myers JC, Jones TA, Pohjolainen E-R, Kadri AS, Goddard AD, Sheer D, Solomon E, and Pihlajaniemi T. Molecular cloning of $\alpha 5(\text{IV})$ collagen and assignment of the gene to the region of the X chromosome containing the Alport syndrome locus. *Am J Hum Genet* 46: 1024-1033, 1990
- A.39 Kivirikko KI, Helaakoski T, Tasanen K, Vuori K, Myllylä R, Parkkonen T, and Pihlajaniemi T. Molecular biology of prolyl 4-hydroxylase. *Ann NY Acad Sci* 580: 132-142, 1990
- A.40 Pihlajaniemi T, Tamminen M, Sandberg M, Hirvonen H, and Vuorio E. The $\alpha 1$ chain of type XIII collagen: polypeptide structure, alternative splicing and tissue distribution. *Ann NY Acad Sci* 580: 440-443, 1990
- A.41 Helaakoski T, Vuori K, Parkkonen T, Myllylä R, Chow LT, Kivirikko KI, and Pihlajaniemi T. Characterization of cDNA and genomic clones for the α subunit of human prolyl 4-hydroxylase. *Ann NY Acad Sci* 580: 473-476, 1990
- A.42 Myllylä R, Pihlajaniemi T, Pajunen L, Turpeenniemi-Hujanen T, and Kivirikko KI. Molecular cloning of chick lysyl hydroxylase. Little homology in primary structure to the two types of subunit of prolyl 4-hydroxylase. *J Biol Chem* 266: 2805-2810, 1991
- A.43 Palosaari PM, Vihinen M, Mäntsälä PI, Alexson SEH, Pihlajaniemi T, and Hiltunen JK. Amino acid sequence similarities of the mitochondrial short-chain Δ^3, Δ^2 -enoyl-CoA isomerase and peroxisomal multifunctional $\Delta 3, \Delta 2$ -enoyl-CoA isomerase, 2-enoyl-CoA hydratase, 3-hydroxyacyl-CoA dehydrogenase enzyme in rat liver. The proposed occurrence of isomerization and hydration in the same catalytic domain of the multifunctional enzyme. *J Biol Chem* 266: 10750-10753, 1991
- A.44 Tikka L, Elomaa O, Pihlajaniemi T, and Tryggvason K. Human $\alpha 1(\text{XIII})$ collagen gene. Multiple forms of the gene transcripts are generated through complex alternative splicing of several short exons. *J Biol Chem* 266: 17713-17719, 1991
- A.45 Autio-Harmainen H, Liakka A, Sandberg M, Pihlajaniemi T, and Vuorio E. Localization of type IV collagen and laminin mRNA in developing human placenta by using in situ hybridization. *Lab Invest* 64: 483-491, 1991
- A.46 Pajunen L, Jones TA, Goddard A, Sheer D, Solomon E, Pihlajaniemi T, and Kivirikko KI. Regional assignment of the human gene coding for a multifunctional polypeptide (P4HB) acting as the β -subunit of prolyl 4-hydroxylase and the enzyme disulphide isomerase to 17q25. *Cytogenet Cell Genet* 56: 165-168, 1991

- A.47 Hämäläinen E-R, Jones TA, Sheer D, Taskinen K, Pihlajaniemi T, and Kivirikko KI. Molecular cloning of human lysyl oxidase and assignment of the gene to chromosome 5q23.3-31.2. *Genomics* 11: 508-516, 1991
- A.48 Boye E, Vetrie D, Flinter F, Buckle B, Pihlajaniemi T, Hämäläinen E- R, Myers JC, Bobrow M, and Harris A. Major rearrangements in the $\alpha 5(\text{IV})$ collagen gene in three patients with Alport syndrome. *Genomics* 11: 1125-1132, 1991
- A.49 Pihlajaniemi T. Concept of one gene one protein revisited: Protein disulphide isomerase, a multifunctional protein. *Ann Med* 23: 393-394, 1991
- A.50 Pihlajaniemi T, Myllylä R, and Kivirikko KI. Prolyl 4-hydroxylase and its role in collagen synthesis. *J Hepatol* 13: 2-7, 1991
- A.51 Vuori K, Pihlajaniemi T, Myllylä R, and Kivirikko KI. Site-directed mutagenesis of human protein disulphide isomerase: effect on the assembly, activity and endoplasmic reticulum retention of human prolyl 4-hydroxylase in *Spodoptera frugiperda* insect cells. *EMBO J* 11: 4213-4217, 1992
- A.52 Vuori K, Pihlajaniemi T, Marttila M, and Kivirikko KI. Characterization of the human prolyl 4-hydroxylase tetramer and its multifunctional protein disulfide-isomerase subunit synthesized in a baculovirus expression system. *Proc Natl Acad Sci USA* 89: 7467-7470, 1992
- A.53 Myers JC, Kivirikko S, Gordon MK, Dion AS, and Pihlajaniemi T. Identification of a previously unknown human collagen chain, $\alpha 1(\text{XV})$, characterized by extensive interruptions in the triple-helical region. *Proc Natl Acad Sci USA* 89: 10144-10148, 1992
- A.54 Vuori K, Myllylä R, Pihlajaniemi T, and Kivirikko KI. Expression and site-directed mutagenesis of human protein disulfide isomerase in *Escherichia coli*. This multifunctional polypeptide has two independently acting catalytic sites for the isomerase activity. *J Biol Chem* 267: 7211-7214, 1992
- A.55 Tasanen K, Oikarinen J, Kivirikko KI, and Pihlajaniemi T. Promoter of the gene for the multifunctional protein disulfide isomerase polypeptide. Functional significance of the six CCAAT boxes and other promoter elements. *J Biol Chem* 267: 11513-11519, 1992
- A.56 Juvonen M, and Pihlajaniemi T. Characterization of the spectrum of alternative splicing of $\alpha 1(\text{XIII})$ collagen transcripts in HT-1080 cells and calvarial tissue resulted in identification of two previously unidentified alternatively spliced sequences, one previously unidentified exon, and nine new mRNA variants. *J Biol Chem* 267: 24693-24699, 1992
- A.57 Juvonen M, Sandberg M, and Pihlajaniemi T. Patterns of expression of the six alternatively spliced exons affecting the structures of the COL1 and NC2 domains of the $\alpha 1(\text{XIII})$ collagen chain in human tissues and cell lines. *J Biol Chem* 267: 24700-24707, 1992
- A.58 Huebner K, Cannizzaro LA, Jabs EW, Kivirikko S, Manzone H, Pihlajaniemi T, and Myers JC. Chromosomal assignment of a gene encoding a new collagen type (COL15A1) to 9q21→q22. *Genomics* 14: 220-224, 1992

- A.59 Renieri A, Seri M, Myers JC, Pihlajaniemi T, Massella L, Rizzoni G, and De Marchi M. *De novo* mutation in the COL4A5 gene converting glycine 325 to glutamic acid in Alport syndrome. *Hum Mol Genet* 1: 127-129, 1992
- A.60 Renieri A, Seri M, Myers JC, Pihlajaniemi T, Sessa A, Rizzoni G, and De Marchi M. Alport syndrome caused by a 5' deletion within the COL4A5 gene. *Hum Genet* 89: 120-121, 1992
- A.61 Tasanen K, Oikarinen J, Kivirikko KI, and Pihlajaniemi T. Interaction of transcription factor Sp1 with the promoter of the gene for the multifunctional protein disulphide isomerase polypeptide. *Biochem J*, 292: 41-45, 1993.
- A.62 Juvonen M, Pihlajaniemi T, and Autio-Harminen H. Location and alternative splicing of type XIII collagen RNA in the early human placenta. *Lab Invest* 69: 541-551, 1993.
- A.63 Hämäläinen E-R, Kempainen R, Pihlajaniemi T, and Kivirikko KI. Structure of the human lysyl oxidase gene. *Genomics* 17: 544-548, 1993.
- A.64 Rehn M, and Pihlajaniemi T. α 1(XVIII), a collagen chain with frequent interruptions in the collagenous sequence, a distinct tissue distribution, and homology with type XV collagen. *Proc Natl Acad Sci USA* 91: 4234-4238, 1994
- A.65 Kivirikko S, Heinämäki P, Rehn M, Honkanen N, Myers JC, and Pihlajaniemi T. Primary structure of the α 1 chain of human type XV collagen and exon-intron organization in the 3' region of the corresponding gene. *J Biol Chem* 269: 4773-4779, 1994
- A.66 Rehn M, Hintikka E, and Pihlajaniemi T. Primary structure of the α 1 chain of mouse type XVIII collagen, partial structure of the corresponding gene, and comparison of the α 1(XVIII) chain with its homologue, the α 1(XV) collagen chain. *J Biol Chem* 269: 13929-13935, 1994
- A.67 Veijola J, Koivunen P, Annunen P, Pihlajaniemi T, and Kivirikko KI. Cloning, baculovirus expression and characterization of the α subunit of prolyl 4-hydroxylase from the nematode *Caenorhabditis elegans*. This α subunit forms an active $\alpha\beta$ dimer with the human protein disulfide isomerase/ β subunit. *J Biol Chem* 269: 26746-26753, 1994
- A.68 Helaakoski T, Veijola J, Vuori K, Rehn M, Chow LT, Taillon-Miller P, Kivirikko KI, and Pihlajaniemi T. Structure and expression of the human gene for the α subunit of prolyl 4-hydroxylase. The two alternatively spliced types of mRNA correspond to two homologous exons the sequences of which are expressed in a variety of tissues. *J Biol Chem* 269: 27847-27853, 1994
- A.69 Helaakoski T, Annunen P, Vuori K, MacNeil IA, Pihlajaniemi T, and Kivirikko KI. Cloning, baculovirus expression, and characterization of a second mouse prolyl 4-hydroxylase α -subunit isoform: Formation of an $\alpha_2\beta_2$ tetramer with the protein disulfide isomerase/ β subunit. *Proc Natl Acad Sci USA* 92:4427-4431, 1995
- A.70 Rehn M, and Pihlajaniemi T. Identification of three N-terminal ends of type XVIII collagen chains and tissue-specific differences in the expression of the corresponding transcripts. The longest form contains a novel motif homologous to rat and *Drosophila* frizzled proteins. *J Biol Chem* 270: 4705-4711, 1995

- A.71 Lamberg A, Pihlajaniemi T, and Kivirikko KI. Site-directed mutagenesis of the α subunit of human prolyl 4-hydroxylase. Identification of three histidine residues critical for catalytic activity. *J Biol Chem* 270: 9926-9931, 1995
- A.72 Hämäläinen E-R, Kemppainen R, Kuivaniemi H, Tromp G, Vaheri A, Pihlajaniemi T, and Kivirikko KI. Quantitative polymerase chain reaction of lysyl oxidase mRNA in malignantly transformed human cell lines demonstrates that their low lysyl oxidase activity is due to low quantities of its mRNA and low levels of transcription of the respective gene. *J Biol Chem* 270: 21590-21593, 1995
- A.73 Kivirikko S, Saarela J, Myers JC, Autio-Harmainen H, and Pihlajaniemi T. Distribution of type XV collagen transcripts in human tissue and their production by muscle cells and fibroblasts. *Am J Pathol* 147: 1500-1509, 1995
- A.74 Pihlajaniemi T, and Rehn M. Two new collagen subgroups: Membrane-associated collagens and types XV and XVIII. *Progr Nucleic Acid Res Mol Biol* 50: 225-262, 1995
- A.75 Pihlajaniemi T. Kollagenfamiljen: Nya medlemmar och nya tankar om deras biosyntes. *Nord Med* 110: 96-99, 1995
- A.76 Lamberg A, Helaakoski T, Myllyharju J, Peltonen S, Notbohm H, Pihlajaniemi T, and Kivirikko KI. Characterization of human type III collagen expressed in a baculovirus system. Production of a protein with a stable triple helix requires coexpression with the two types of recombinant prolyl 4-hydroxylase subunit. *J Biol Chem* 271: 11988-11995, 1996
- A.77 Mazzorana M, Snellman A, Kivirikko KI, van der Rest, and Pihlajaniemi T. Involvement of prolyl 4-hydroxylase in the assembly of trimeric minicollagen XII. Study in a baculovirus expression system. *J Biol Chem* 271: 29003-29008, 1996
- A.78 Rehn M, Hintikka E, and Pihlajaniemi, T. Characterization of the mouse gene for the $\alpha 1$ chain of type XVIII collagen (*Col18a1*) reveals that the three variant N-terminal polypeptide forms are transcribed from two widely separated promoters. *Genomics* 32: 436-446, 1996
- A.79 Lamberg A, Jauhiainen M, Metso J, Ehnholm C, Shoulders C, Scott J, Pihlajaniemi T, and Kivirikko KI. The role of protein disulphide isomerase in the microsomal triacylglycerol transfer protein does not reside in its isomerase activity. *Biochem J* 315: 533-536, 1996
- A.80 Veijola J, Pihlajaniemi T, and Kivirikko, K.I. Co-expression of the α subunit of human prolyl 4-hydroxylase with BiP polypeptide in insect cells leads to the formation of soluble and insoluble complexes. Soluble α -subunit-BiP complexes have no prolyl 4-hydroxylase activity. *Biochem J* 315: 613-618, 1996
- A.81 Koivunen P, Helaakoski T, Annunen P, Veijola J, Räisänen S, Pihlajaniemi T, and Kivirikko, K.I. ERp60 does not substitute for protein disulphide isomerase as the β subunit of prolyl 4-hydroxylase. *Biochem J* 316: 599-605, 1996
- A.82 Veijola J, Annunen P, Koivunen P, Page A, Pihlajaniemi T, and Kivirikko KI. Baculovirus expression of two protein disulphide isomerase isoforms from *Caenorhabditis elegans* and characterization of prolyl 4-hydroxylases containing one of these polypeptides as their β subunit. *Biochem J* 317: 721-729, 1996

- A.83 Kemppainen R, Hämäläinen E-R, Kuivaniemi H, Tromp G, Pihlajaniemi T, and Kivirikko KI. Expression of mRNAs for lysyl oxidase and type III procollagen in cultured fibroblasts from patients with the Menkes and occipital horn syndromes as determined by quantitative polymerase chain reaction. *Arch Biochem Biophys* 328: 101-106, 1996
- A.84 Hämäläinen E-R, Renieri A, Pecoraro C, De Marchi M, and Pihlajaniemi T. Unequal homologous crossing over resulting in duplication of 36 base pairs within exon 47 of the COL4A5 gene in a family with Alport syndrome. *Hum Mut* 8: 265-269, 1996
- A.85 Rehn M, and Pihlajaniemi T. Type XV and XVIII collagens, a new subgroup among the family of collagens. *Semin Cell Biol* 7: 673-679, 1996
- A.86 Vuorela A, Myllyharju J, Nissi R, Pihlajaniemi T, and Kivirikko KI. Assembly of human prolyl 4-hydroxylase and type III collagen in the yeast *Pichia pastoris*: formation of a stable enzyme tetramer requires coexpression with collagen and assembly of a stable collagen requires coexpression with prolyl 4-hydroxylase. *EMBO J* 16: 6702-6712, 1997
- A.87 Annunen P, Helaakoski T, Myllyharju J, Veijola J, Pihlajaniemi T, and Kivirikko KI. Cloning of the human prolyl 4-hydroxylase α subunit isoform α (II) and characterization of the type II enzyme tetramer. The α (I) and α (II) subunits do not form a mixed α (I) α (II) β ₂ tetramer. *J Biol Chem* 272: 17342-17348, 1997
- A.88 Myllyharju J, Lamberg A, Notbohm H, Fietzek PP, Pihlajaniemi T, and Kivirikko KI. Expression of wild-type and modified pro α chains of human type I procollagen in insect cells leads to the formation of stable [α 1(I)]₂ α 2(I) collagen heterotrimers and [α 1(I)]₃ homotrimers but not [α 2(I)]₃ homotrimers. *J Biol Chem* 272: 21824-21830, 1997
- A.89 Hägg PM, Hägg PO, Peltonen S, Autio-Harmainen H, and Pihlajaniemi T. Location of type XV collagen in human tissues and its accumulation in the interstitial matrix of the fibrotic kidney. *Am J Pathol* 150: 2075-2086, 1997
- A.90 Hägg PM, Horelli-Kuitunen N, Eklund L, Palotie A, and Pihlajaniemi T. Cloning of mouse type XV collagen sequences and mapping of the corresponding gene to 4B1-3. Comparison of mouse and human α 1(XV) collagen sequences indicates divergence in the number of small collagenous domains. *Genomics* 45: 31-41, 1997
- A.91 Horelli-Kuitunen N, Kvist A-P, Helaakoski T, Kivirikko KI, Pihlajaniemi T, and Palotie A. The order and transcriptional orientation of the human COL13A1 and P4HA genes on chromosome 10 long arm determined by high-resolution FISH. *Genomics* 46: 299-302, 1997
- A.92 Peltonen S, Rehn M, and Pihlajaniemi T. Alternative splicing of mouse α 1(XIII) collagen RNAs results in at least 17 different transcripts, predicting α 1(XIII) collagen chains with length varying between 651 and 710 amino acid residues. *DNA Cell Biol* 16: 227-234, 1997
- A.93 Hägg P, Rehn M, Huhtala P, Väisänen T, Tamminen M, and Pihlajaniemi T. Type XIII collagen is identified as a plasma membrane protein. *J Biol Chem* 273: 15590-15597, 1998

- A.94 Hägg PM, Muona A, Liétard J, Kivirikko S, and Pihlajaniemi T. Complete exon-intron organization of the human gene for the $\alpha 1$ chain of type XV collagen (COL15A1) and comparison with the homologous *Coll8a1* gene. *J Biol Chem* 273: 17824-17831, 1998
- A.95 Saarela J, Rehn M, Oikarinen A, Autio-Harmainen H, and Pihlajaniemi T. The short and long forms of type XVIII collagen show clear tissue specificities in their expression and location in basement membrane zones in man. *Am J Pathol* 153: 611-626, 1998
- A.96 Musso O, Rehn M, Saarela J, Th  ret N, Li  tard J, Hintikka E, Lotrian D, Campion J-P, Pihlajaniemi T, and Cl  ment B. Collagen XVIII is localized in sinusoids and basement membrane zones and expressed by hepatocytes and activated stellate cells in fibrotic human liver. *Hepatology* 28: 98-107, 1998
- A.97 Saarela J, Ylik  rpp   R, Rehn M, Purmonen S, and Pihlajaniemi T. Complete primary structure of two variant forms of human type XVIII collagen and tissue-specific differences in the expression of the corresponding transcripts. *Matrix Biol* 16: 319-328, 1998
- A.98 Nokelainen M, Helaakoski T, Myllyharju J, Notbohm H, Pihlajaniemi T, Fietzek PP, and Kivirikko KI. Expression and characterization of recombinant human type II collagens with low and high contents of hydroxylysine and its glycosylated forms. *Matrix Biol* 16: 329-338, 1998
- A.99 Kivirikko KI, and Pihlajaniemi T. Collagen hydroxylases and the protein disulfide isomerase subunit of prolyl 4-hydroxylases. *Adv Enzymol Related Areas Mol Biol* 72: 325-398, 1998
- A.100 Rehn M, Pihlajaniemi T, Hofmann K, and Bucher P. The frizzled motif: in how many different protein families does it occur? *Trends Biochem Sci* 23: 415-417, 1998
- A.101 Myers LK, Brand DD, Ye XJ, Cremer MA, Rosloniec EF, Bodo M, Myllyharju J, Helaakoski T, Nokelainen M, Pihlajaniemi T, Kivirikko KI, Yang CL, Ala-Kokko L, Prockop DJ, Notbohm H, Fietzek P, Stuart JM, and Kang AH. Characterization of recombinant type II collagen: arthritogenicity and tolerogenicity in DBA/1 mice. *Immunol* 95: 631-639, 1998
- A.102 Valtola R, Salven P, Heikkil   P, Taipale J, Joensuu H, Rehn M, Pihlajaniemi T, Weich H, deWaal R, and Alitalo K. VEGFR-3 and its ligand VEGF-C are associated with angiogenesis in breast cancer. *Am J Pathol* 154: 1381-1390, 1999
- A.103 Kvist A-P, Latvanlehto A, Sund M, Horelli-Kuitunen N, Rehn M, Palotie A, Beier D, and Pihlajaniemi T. Complete exon-intron organization and chromosomal location of the gene for mouse type XIII collagen (col13a1) and comparison with its human homologue. *Matrix Biol* 18: 261-274, 1999
- A.104 Kivirikko S, Mauviel A, Pihlajaniemi T, and Uitto J. Cytokine modulation of type XV collagen gene expression in human dermal fibroblast cultures. *Exp Dermatol* 8: 407-412, 1999
- A.105 Peltonen S, Hentula M, H  gg P, Yl  -Outinen H, Tuukkanen J, Lakkakorpi J, Rehn M, Pihlajaniemi T, and Peltonen J. A novel component of epidermal cell-matrix and cell-cell contacts: transmembrane protein type XIII collagen. *J Invest Dermatol* 113: 635-642, 1999

- A.106 Linnola RJ, Sund M, Ylönen R, and Pihlajaniemi T. Adhesion of soluble fibronectin, laminin, and collagen type IV to intraocular lens materials. *J Cataract Refract Surg* 25: 1486-1491, 1999
- A.107 Vuorela A, Myllyharju J, Pihlajaniemi T, and Kivirikko KI. Coexpression with collagen markedly increases the half-life of the recombinant human prolyl 4-hydroxylase tetramer in the yeast *Pichia pastoris*. *Matrix Biol* 18: 519-522, 1999
- A.108 Snellman A, Tu H, Väisänen T, Kvist A-P, Huhtala P, and Pihlajaniemi T. A short sequence in the N-terminal region is required for the trimerization of type XIII collagen and is conserved in other collagenous transmembrane proteins. *EMBO J* 19: 5051-5059, 2000
- A.109 Nykvist P, Tu H, Ivaska J, Käpylä J, Pihlajaniemi T, and Heino J. Distinct recognition of collagen subtypes by $\alpha 1\beta 1$ and $\alpha 2\beta 1$ integrins - $\alpha 1\beta 1$ mediates cell adhesion to type XIII collagen. *J Biol Chem* 275: 8255-8261, 2000
- A.110 Snellman A, Keränen M-R, Hägg PO, Lamberg A, Hiltunen JK, Kivirikko KI, and Pihlajaniemi T. Type XIII collagen forms homotrimers with three triple-helical collagenous domains and its association into disulfide-bonded trimers is enhanced by prolyl 4-hydroxylase. *J Biol Chem* 275: 8936-8944, 2000.
- A.111 Liétard J, Théret N, Rehn M, Musso O, Dargère D, Pihlajaniemi T, and Clément B. The promoter of the long variant of collagen XVIII, the precursor of endostatin, contains liverspecific regulatory elements. *Hepatology*, 32:1377-1385, 2000
- A.112 Wagner K, Pöschl E, Turnay J, Baik J, Pihlajaniemi T, Frischholz S, and van der Mark K. Coexpression of α and β subunits of prolyl 4-hydroxylase stabilizes the triple helix of recombinant human type X collagen. *Biochem J* 352:907-911, 2000
- A.113 Keizer-Gunnink I, Vuorela A, Myllyharju J, Pihlajaniemi T, Kivirikko KI, and Veenhuis M. Accumulation of properly folded human type III procollagen molecules in specific intracellular membraneous compartments in the yeast *Pichia pastoris*. *Matrix Biol* 19: 29-36, 2000
- A.114 Eklund L, Muona A, Liétard J, and Pihlajaniemi T. Structure of the mouse type XV collagen gene, *Col15a1*, comparison with the human *COL15A1* gene and functional analysis of the promoters of both genes. *Matrix Biol* 19: 489-500, 2000
- A.115 Sandberg-Lall M, Hägg PO, Wahlström I, and Pihlajaniemi T. Type XIII collagen is widely expressed in the adult and developing human eye and accentuated in the ciliary muscle, the optic nerve and the neural retina. *Exp Eye Res* 70: 401-410, 2000
- A.116 Sund M, Ylönen R, Tuomisto A, Sormunen R, Tahkola J, Kvist A-P, Kontusaari S, Autio-Harmainen H, and Pihlajaniemi T. Abnormal adherence junctions in the heart and reduced angiogenesis in transgenic mice overexpressing mutant type XIII collagen. *EMBO J* 20: 5153-5164, 2001
- A.117 Rehn M, Veikkola T, Kukk-Valdre E, Nakamura H, Ilmonen M, Lombardo CL, Pihlajaniemi T, Alitalo K, and Vuori K. Interaction of endostatin with integrins implicated in angiogenesis. *Proc Natl Acad Sci USA* 98: 1024-1029, 2001

- A.118 Eklund L, Piuhola J, Komulainen J, Sormunen R, Ongvarrasopone C, Fässler R, Muona A, Ilves M, Ruskoaho H, Takala T, and Pihlajaniemi T. Lack of type XV collagen causes a skeletal myopathy and cardiovascular defects in mice. *Proc Natl Acad Sci USA* 98: 1194-1199, 2001
- A.119 Ackley B, Crew JR, Elamaa H, Pihlajaniemi T, Kuo CJ, and Kramer JM. The NC1/endostatin domain of *C. elegans* type XVIII collagen affects cell migration and axon guidance. *J Cell Biol* 152: 1219-1232, 2001
- A.120 Lin Y, Zhang S, Rehn M, Itäranta P, Tuukkanen J, Heljasvaara R, Peltoketo H, Pihlajaniemi T, and Vainio S. Induced repatterning of type XVIII collagen expression in ureter bud from kidney to lung type: association with sonic hedgehog and ectopic surfactant protein C. *Development* 128: 1573-1585, 2001
- A.121 Musso O, Rehn M, Théret N, Turlin B, Bioulac-Sage P, Lotrian D, Champion J-P, Pihlajaniemi T, and Clément B. Tumor progression is associated with significant decrease in the expression of the endostatin precursor collagen XVIII in human hepatocellular carcinomas. *Cancer Res* 61: 45-49, 2001
- A.122 Wickström SA, Veikkola T, Rehn M, Pihlajaniemi T, Alitalo K, and Keski-Oja, J. Endostatin-induced modulation of plasminogen activation with concomitant loss of focal adhesions and actin stress fibers in cultured human endothelial cells. *Cancer Res*, 61: 6511-6516, 2001
- A.123 Kvist A-P, Latvanlehto A, Sund M, Eklund L, Väisänen T, Hägg PO, Sormunen R, Komulainen J, Fässler R, and Pihlajaniemi T. Lack of cytosolic and transmembrane domains of type XIII collagen results in progressive myopathy. *Am J Pathol* 159: 1581-1592, 2001
- A.124 Musso O, Theret N, Heljasvaara R, Rehn M, Turlin B, Champion JP, Pihlajaniemi T, and Clement B. Tumor hepatocytes and basement membrane-producing cells specifically express two different forms of the endostatin precursor, collagen XVIII, in human liver cancers. *Hepatology* 33: 868-876, 2001
- A.125 Hägg PO, Väisänen T, Tuomisto A, Rehn M, Tu H, Huhtala P, Eskelinen S, and Pihlajaniemi T. Type XIII collagen: a novel focal adhesion component present in a range of cell-matrix adhesions and in the intercalated discs between cardiac muscle cells. *Matrix Biol* 19:727-742, 2001
- A.126 Sund M, Väisänen T, Kaukinen S, Ilves M, Tu H, Autio-Harmainen H, Rauvala H, and Pihlajaniemi T. Distinct expression of type XIII collagen in neuronal structures and other tissues during mouse development. *Matrix Biol* 20: 215-231, 2001
- A.127 Sallinen R, Latvanlehto A, Kvist A-P, Rehn M, Eerola I, Chu M-L, Bonaldo P, Saitta B, Bressan GM, Pihlajaniemi T, Vuorio E, Palotie A, Wessman M, and Horelli-Kuitunen N. Physical mapping of mouse collagen genes on chromosome 10 by high-resolution FISH. *Mamm Genome* 12: 340-346, 2001
- A.128 *Fukai N, *Eklund L, *Marneros AG, Oh SP, Keene DR, Tamarkin L, Niemelä M, Ilves M, Li E, Pihlajaniemi T, and Olsen BR. Lack of collagen XVIII/endostatin results in eye abnormalities. *EMBO J* 21: 1535-1544, 2002. *Equal contribution

- A.129 Tu H, Sasaki T, Snellman A, Göhring W, Pirilä P, Timpl R, and Pihlajaniemi T. The type XIII collagen ectodomain is a 150-nm rod and capable of binding to fibronectin, nidogen-2, perlecan and heparin. *J Biol Chem* 277:23092-23099, 2002
- A.130 Van Horssen J, Wilhelmus MMM, Heljasvaara R, Pihlajaniemi T, Wesseling P, de Waal RMW, and Verbeek MM. Collagen XVIII: a novel heparan sulfate proteoglycan associated with vascular amyloid depositions and senile plaques in Alzheimer's disease brains. *Brain Pathol* 12: 456-462, 2002
- A.131 Muona A, Eklund L, Väisänen T, and Pihlajaniemi T. Developmentally regulated expression of type XV collagen correlates with abnormalities in *Coll15a1*^{-/-} mice. *Matrix Biol* 21: 89-102, 2002
- A.132 Elamaa H, Peterson J, Pihlajaniemi T, and Destrée O. Cloning of three variants of type XVIII collagen and their expression patterns during *Xenopus laevis* development. *Mech Dev* 114: 109-113, 2002
- A.133 Kawashima H, Watanabe N, Hirose M, Sun X, Atarashi K, Kimura T, Shikata K, Matsuda M, Ogawa D, Heljasvaara R, Rehn M, Pihlajaniemi T, and Miyasaka M. Collagen XVIII, a basement membrane heparan sulfate proteoglycan, interacts with L-selectin and monocyte chemoattractant protein-1. *J Biol Chem* 278: 13069-13076, 2003
- A.134 Nyberg P, Heikkilä P, Sorsa T, Luostarinen J, Heljasvaara R, Stenman UH, Pihlajaniemi T, and Salo T. Endostatin inhibits human tongue carcinoma cell invasion and intravasation and blocks the activation of matrix metalloprotease-2, -9 and -13. *J Biol Chem* 278: 22404-22411, 2003
- A.135 Latvanlehto A, Snellman A, Tu H, and Pihlajaniemi T. Type XIII collagen and some other transmembrane collagens contain two separate coiled-coil motifs, which may function as independent oligomerization domains. *J Biol Chem* 278: 37590-37599, 2003
- A.136 Ylikärppä R, Eklund L, Sormunen R, Kontiola AI, Utriainen A, Määttä M, Fukai N, Olsen BR, and Pihlajaniemi T. Lack of type XVIII collagen results in anterior ocular defects. *FASEB J* 17: 2257-2259, 2003
- A.137 Vainio S, Lin Y, and Pihlajaniemi T. Induced repatterning of type XVIII collagen associates with ectopic sonic hedgehog and lung surfactant C gene expression and changes in epithelial epigenesis in the ureteric bud. *J Am Soc Nephrol* 14: S3-8, 2003
- A.138 Linnola RJ, Sund M, Ylönen R, and Pihlajaniemi T. Adhesion of soluble fibronectin, vitronectin, and collagen type IV to intraocular lens materials. *J Cataract Refract Surg* 29: 146-152, 2003
- A.139 Lin Y, Zhang S, Tuukkanen J, Peltoketo H, Pihlajaniemi T, and Vainio S. Patterning parameters associated with the branching of the ureteric bud regulated by epithelial-mesenchymal interactions. *Int J Dev Biol* 47: 3-13, 2003
- A.140 Elamaa H, Snellman A, Rehn M, Autio-Harmainen H, and Pihlajaniemi T. Characterization of the human type XVIII collagen gene structure and proteolytic processing and tissue location of the variant containing a frizzled motif. *Matrix Biol*, 22, 427-442, 2003

- A.141 Ylikärppä R, Eklund L, Sormunen R, Muona A, Fukai N, Olsen BR, and Pihlajaniemi T. Double knockout mice reveal a lack of a major functional compensation between collagens XV and XVIII. *Matrix Biol* 22: 443-448, 2003
- A.142 Utriainen A, Sormunen R, Kettunen M, Carvalhaes LS, Sajanti E, Eklund L, Kauppinen R, Kitten GT, and Pihlajaniemi T. Structurally altered basement membranes and hydrocephalus in a type XVIII collagen deficient mouse line. *Hum Mol Genet* 13: 2089-2099, 2004
- A.143 Väisänen M-R, Väisänen T, and Pihlajaniemi T. The shed ectodomain of type XIII collagen affects cell behaviour in matrix-dependent manner. *Biochem J* 380: 685-693, 2004
- A.144 Väänänen A, Tjäderhane L, Eklund L, Heljasvaara R, Pihlajaniemi T, Herva R, Ding Y, Bartlett J, and Salo T. Expression of collagen XVIII and MMP-20 in developing tooth and odontogenic tumors. *Matrix Biol* 23: 153-161, 2004
- A.145 Elamaa H, Sormunen R, Rehn M, Soininen R, and Pihlajaniemi T. Endostatin overexpression specifically in the lens and skin leads to cataract and ultrastructural alterations in basement membranes. *Am J Pathol* 166: 221-229, 2005
- A.146 Ylönen R, Kyrölahti T, Sund M, Ilves M, Lehenkari P, Tuukkanen J, and Pihlajaniemi T. Type XIII collagen strongly affects bone formation in transgenic mice. *J Bone Miner Res* 20: 1381-1393, 2005
- A.147 Hurskainen M, Eklund L, Hägg PO, Fruttiger M, Sormunen R, Ilves M, and Pihlajaniemi T. Abnormal maturation of the retinal vasculature in type XVIII collagen/endostatin deficient mice and changes in retinal glial cells due to lack of collagen types XV and XVIII. *FASEB J* 19: 1564-1566, 2005
- A.148 Heljasvaara R, Nyberg P, Luostarinen J, Parikka M, Heikkilä P, Rehn M, Sorsa T, Salo T, and Pihlajaniemi T. Generation of biologically active endostatin fragments from human collagen XVIII by distinct matrix metalloproteases. *Exp Cell Res* 307: 292-304, 2005
- A.149 Väisänen T, Väisänen M-R, Autio-Harmainen H, and Pihlajaniemi T. Type XIII collagen expression is induced during malignant transformation in various epithelial and mesenchymal tumours. *J Pathol* 207: 324-335, 2005
- A.150 Määttä M, Butzow R, Luostarinen J, Petäjäniemi N, Pihlajaniemi T, Salo S, Miyazaki K, Autio-Harmainen H, and Virtanen I. Differential expression of laminin isoforms in ovarian epithelial carcinomas suggesting different origin and providing tools for differential diagnosis. *J Histochem Cytochem* 53: 1293-1300, 2005
- A.151 Väisänen T, Väisänen MR, and Pihlajaniemi T. Modulation of the cellular cholesterol level affects shedding of the type XIII collagen ectodomain. *J Biol Chem* 281: 33352-33362, 2006
- A.152 Väisänen M-R, Väisänen T, Tu H, Pirilä P, Sormunen R, and Pihlajaniemi T. The shed ectodomain of type XIII collagen associates with the fibrillar matrix and may interfere with its assembly in vitro. *Biochem J* 393: 43-50, 2006
- A.153 Carvalhaes LS, Gervásio OL, Guatimosim C, Heljasvaara R, Sormunen R, Pihlajaniemi T, and Kitten GT. Collagen XVIII/endostatin is associated with the epithelial-

mesenchymal transformation in the atrioventricular valves during cardiac development. *Dev Dyn*, 235: 132-142, 2006

- A.154 Määttä M, Väisänen T, Väisänen M-R, Pihlajaniemi T, and Tervo T. Altered expression of type XIII collagen in keratoconus and scarred human cornea: Increased expression in scarred cornea is associated with myofibroblast transformation. *Cornea* 25: 448-453, 2006
- A.155 Määttä M, Heljasvaara R, Sormunen R, Pihlajaniemi T, Autio-Harmainen H, and Tervo T. Differential expression of collagen types XVIII/endostatin and XV in normal, keratoconus, and scarred human corneas. *Cornea* 25: 341-349, 2006
- A.156 Snellman A, Tuomisto A, Koski A, Latvanlehto A, and Pihlajaniemi T. The role of disulfide bonds and alpha-helical coiled-coils in the biosynthesis of type XIII collagen and other collagenous transmembrane proteins. *J Biol Chem* 282: 14898-14905, 2007
- A.157 Brideau G, Mäkinen MJ, Elamaa H, Alitalo K, and Pihlajaniemi T, Heljasvaara R. Endostatin overexpression inhibits lymphangiogenesis and lymph node metastasis in mice. *Cancer Research* 67: 11528-11535, 2007
- A.158 Celie JW, Rutjes NW, Keuning ED, Soinen R, Heljasvaara R, Pihlajaniemi T, Dräger AM, Zweegman S, Kessler FL, Beelen RH, Florquin S, Aten J, and van den Born J. Subendothelial heparan sulfate proteoglycans become major L-selectin and monocyte chemoattractant protein-1 ligands upon renal ischemia/reperfusion. *Am J Pathol* 170: 1865-1878, 2007
- A.159 Vainionpää N, Bützow R, Hukkanen M, Jackson DG, Pihlajaniemi T, Sakai LY, and Virtanen I. Basement membrane protein distribution in LYVE-1-immunoreactive lymphatic vessels of normal tissues and ovarian carcinomas. *Cell Tissue Res* 328: 317-28, 2007
- A.160 Määttä M, Heljasvaara R, Pihlajaniemi T, and Uusitalo M. Collagen XVIII/endostatin shows a ubiquitous distribution in human ocular tissues and endostatin-containing fragments accumulate in ocular fluid samples. *Graefes Arch Clin Exp Ophthalmol* 245: 74-81, 2007
- A.161 Quélard D, Lavergne E, Hendaoui I, Elamaa H, Tiirola U, Heljasvaara R, Pihlajaniemi T, Clément B, and Musso O. A cryptic frizzled module in cell surface collagen 18 inhibits Wnt/beta-catenin signaling. *PLoS One* 3: e1878, 2008
- A.162 Tuomisto A, Sund M, Tahkola J, Latvanlehto A, Savolainen E-R, Autio-Harmainen H, Liakka A, Sormunen R, Vuoristo J, West A, Lahesmaa R, Morse HC 3rd, and Pihlajaniemi T. A mutant collagen XIII alters intestinal expression of immune response genes and predisposes transgenic mice to develop B-cell lymphomas. *Cancer Res* 68: 10324-10332, 2008
- A.163 Seppinen L, Sormunen R, Soini Y, Elamaa H, Heljasvaara R, and Pihlajaniemi T. Lack of collagen XVIII accelerates cutaneous wound healing, while overexpression of its endostatin domain leads to delayed healing. *Matrix Biol*, 27: 535-546, 2008
- A.164 Tahkola J, Räsänen J, Sund M, Mäkikallio K, Autio-Harmainen H, and Pihlajaniemi T. Cardiac dysfunction in transgenic mouse fetuses overexpressing shortened type XIII collagen. *Cell Tissue Res* 333: 61-69, 2008

- A.165 Perkins GD, Nathani N, Richter AG, Park D, Shyamsundar M, Heljasvaara R, Pihlajaniemi T, Manji M, Tunnicliffe W, McAuley D, Gao F, and Thickett DR Type XVIII collagen degradation products in acute lung injury. *Crit Care* 13:R52, 2009
- A.166 Suzuki O, Kague E, Bagatini K, Tu H, Heljasvaara R, Carvalhaes L, Gava E, de Oliveira G, Godoi P, Oliva G, Kitten G, Pihlajaniemi T, and Passos-Bueno MR. Novel pathogenic mutations and skin biopsy analysis in Knobloch syndrome. *Mol Vis* 15: 801-809, 2009
- A.167 Huilaja L, Hurskainen T, Autio-Harmanen H, Sormunen R, Tu H, Hofmann SC, Pihlajaniemi T, Bruckner-Tuderman L, and Tasanen K. Glycine substitution mutations cause intracellular accumulation of collagen XVII and affect its post-translational modifications. *J Invest Dermatol* 129: 2302–2306, 2009
- A.168 Sipola A, Seppinen L, Pihlajaniemi T, and Tuukkanen J. Endostatin affects osteoblast behavior in vitro, but collagen XVIII/endostatin is not essential for skeletal development in vivo. *Calcif Tissue Int* 85: 412-420, 2009
- A.169 Thulin Å, Ringvall M, Dimberg A, Kårehed K, Väisänen T, Väisänen M-R, Hamad O, Wang J, Bjerkvig R, Nilsson B, Pihlajaniemi T, Åkerud H, Pietras K, Jahnen-Dechent W, Siegbahn A, and Olsson A-K. Activated platelets provide a functional microenvironment for the anti-angiogenic fragment of histidine-rich glycoprotein. *Mol Cancer Res*, 7: 1792-1802, 2009
- A.170 Hurskainen M, Ruggiero F, Hägg P, Pihlajaniemi T, and Huhtala P. Recombinant human collagen XV regulates cell adhesion and migration. *J Biol Chem* 285: 5258-5265, 2010
- A.171 Bruckner-Tuderman L, von der Mark K, Pihlajaniemi T, and Unsicker K. Cell interactions with the extracellular matrix. *Cell Tissue Res* 339: 1-5, 2010
- A.172 Latvanlehto A, Fox MA, Sormunen R, Tu H, Oikarainen T, Koski A, Naumenko N, Shakirzyanova A, Kallio M, Ilves M, Giniatullin R, Sanes JR, and Pihlajaniemi T. Muscle-derived collagen XIII regulates maturation of the skeletal neuromuscular junction. *J Neurosci* 30: 12230-12241, 2010
- A.173 Rasi K, Hurskainen M, Kallio M, Stavén S, Sormunen R, Heape AM, Avila RL, Kirschner D, Muona A, Tolonen U, Tanila H, Huhtala P, Soininen R, and Pihlajaniemi T. Lack of collagen XV impairs peripheral nerve maturation and, when combined with laminin-411 deficiency, leads to basement membrane abnormalities and sensorimotor dysfunction. *J Neurosci* 30: 14490-14501, 2010
- A.174 Dennis J, Meehan DT, Delimont D, Zallocchi M, Perry GA, O'Brien S, Tu H, Pihlajaniemi T, and Cosgrove D. Collagen XIII induced in vascular endothelium mediates $\alpha\beta 1$ integrin-dependent transmigration of monocytes in renal fibrosis. *Am J Pathol* 177: 2527-2540, 2010
- A.175 Rasi K, Piuhola J, Czabanka M, Sormunen R, Ilves M, Leskinen H, Rysä J, Kerkelä R, Janmey P, Heljasvaara R, Peuhkurinen K, Vuolteenaho O, Ruskoaho H, Vajkoczy P, Pihlajaniemi T, and Eklund L. Collagen XV is necessary for modeling of the extracellular matrix and its deficiency predisposes to cardiomyopathy. *Circ Res* 107: 1241-1252, 2010
- A.176 Seppinen L, and Pihlajaniemi T. The multiple functions of collagen XVIII in development and disease. *Matrix Biol* 30: 83-92, 2011

- A.177 Kinnunen AI, Sormunen R, Elamaa H, Seppinen L, Miller RT, Ninomiya Y, Janmey PA, and Pihlajaniemi T. Lack of collagen XVIII long isoforms affects kidney podocytes, whereas the short form is needed in the proximal tubular basement membrane. *J Biol Chem* 286: 7755-7764, 2011
- A.178 Farshchian M, Kivisaari A, Ala-aho R, Riihilä P, Kallajoki M, Grénman R, Peltonen J, Pihlajaniemi T, Heljasvaara R, and Kähäri V-M. Serpin peptidase inhibitor clade A member 1 (Serpina1) is a novel biomarker for progression of cutaneous squamous cell carcinoma. *Am J Pathol* 179: 1110-1119, 2011
- A.179 Holopainen T, López-Alpuche V*, Zheng W*, Heljasvaara R, Jones D, He Y, Tvorogov D, D'Amico G, Wiener Z, Andersson LC, Pihlajaniemi T, Min W and Alitalo K. Deletion of the endothelial Bmx tyrosine kinase decreases tumor angiogenesis and growth. *Equal contribution. *Cancer Res* 72: 3512-3521, 2012
- A.180 Heikkinen A, Tu H, and Pihlajaniemi T. Collagen XIII: a type II transmembrane protein with relevance to musculoskeletal tissues, microvessels and inflammation. *Int J Biochem & Cell Biol* 44: 714-717, 2012
- A.181 Su J, Stenbjorn RS, Gorse K, Su K, Hauser KF, Ricard-Blum S, Pihlajaniemi T, Fox MA. Target-derived matricryptins organize cerebellar synapse formation through $\alpha 3\beta 1$ integrins. *Cell Rep* 2: 223-230, 2012
- A.182 Aikio M, Alahuhta I, Nurmenniemi S, Suojanen J, Palovuori R, Teppo S, Sorsa T, López-Otín C, Pihlajaniemi T, Salo T, Heljasvaara R, and Nyberg P. Arresten, a collagen-derived angiogenesis inhibitor suppresses oral carcinoma invasiveness. *PLoS ONE* 7: e51044, 2012
- A.183 Field-Ridley A, Heljasvaara R, Pihlajaniemi T, Adatia I, Sun C, Keller RL, Gong WH, Datar S, Oishi P, and Fineman JR. Endostatin, an inhibitor of angiogenesis, decreases after bidirectional superior cavopulmonary anastomosis. *Pediatr Cardiol* 34: 291-295, 2013
- A.184 Salokorpi N, Yrjänä S, Tuominen H, Karttunen A, Heljasvaara R, Pihlajaniemi T, Heikkinen E, and Koivukangas J. Expression of VEGF and collagen XVIII in meningiomas: correlations with histopathological and MRI characteristics. *Acta Neurochir (Wien)*, 155: 989-996, 2013
- A.185 Haikarainen T, Koivunen J, Narwal M, Venkannagari H, Obaji E, Joensuu P, Pihlajaniemi T, and Lehtiö L. Para substituted 2-phenyl-3,4-dihydroquinazolin-4-ones as potent and selective tankyrase inhibitors. *ChemMedChem*, 8: 1978-1985, 2013
- A.186 Aikio M, Hurskainen M, Brideau G, Hägg H, Sormunen R, Heljasvaara R, Gould DB, and Pihlajaniemi T. Collagen XVIII short isoform is critical for retinal vascularization, and overexpression of the Tsp-1 domain affects eye growth and cataract formation. *Invest Ophthalmol Vis Sci* 54: 7450-7462, 2013
- A.187 Narwal M, Koivunen J, Haikarainen T, Venkannagari H, Obaji E, Joensuu P, Pihlajaniemi T, and Lehtiö L. Discovery of Tankyrase Inhibiting Flavones with Increased Potency and Isoenzyme Selectivity. *J Med Chem* 56: 7880-7889, 2013

- A.188 Rygh CB, Løkka G, Heljasvaara R, Taxt T, Pavlin T, Sormunen R, Pihlajaniemi T, Curry FR, Tenstad O, Reed RK. Image-based assessment of microvascular function and structure in Collagen XV and XVIII deficient mice. *J Physiol* 592: 325-336, 2014
- A.189 Aikio M, Elamaa H, Vicente D, Izzi V, Kaur I, Seppinen L, Speedy HE, Kaminska D, Kuusisto S, Sormunen R, Heljasvaara R, Jones EL, Muilu M, Jauhiainen M, Pihlajamäki J, Savolainen MJ, Shoulders CC, and Pihlajaniemi T. Specific collagen XVIII isoforms promote adipose tissue accrual via mechanisms determining adipocyte number and affect fat deposition. *Proc Natl Acad Sci USA* 111: E3043-3052, 2014
- A.190 Zaferani A, Talsma DT, Yazdani S, Celie JW, Aikio M, Heljasvaara R, Navis GJ, Pihlajaniemi T, and van den Born J. Basement membrane zone collagens XV and XVIII/proteoglycans mediate leukocyte influx in renal ischemia/reperfusion. *Plos One* 9: e106732, 2014
- A.191 Kantola T, Väyrynen JP, Klintrup K, Mäkelä J, Karppinen SM, Pihlajaniemi T, Autio-Harmainen H, Karttunen TJ, Mäkinen MJ, and Tuomisto A. Serum endostatin levels are elevated in colorectal cancer and correlate with invasion and systemic inflammatory markers. *Brit J Cancer*, 111: 1605-1613, 2014
- A.192 Heikkinen A, Pihlajaniemi T, Faissner A, and Yuzaki M. Neural ECM and synaptogenesis. *Prog Brain Res* 214: 29-51, 2014
- A.193 Farshchian M, Nissinen L, Siljamäki E, Riihilä P, Toriseva M, Kivisaari A, Ala-aho R, Kallajoki M, Veräjänkorva E, Honkanen HK, Heljasvaara R, Pihlajaniemi T, Grénman R, Peltonen J, Peltonen S, and Kähäri VM. EphB2 promotes progression of cutaneous squamous cell carcinoma. *J Invest Dermatol* 135: 1882-1892, 2015
- A.194 Alahuhta I, Aikio M, Väyrynen O, Nurmenniemi S, Suojanen J, Teppo S, Pihlajaniemi T, Heljasvaara R, Salo T, and Nyberg P. Endostatin induces proliferation of oral carcinoma cells but its effect on invasion is modified by the tumor microenvironment. *Exp Cell Res* 336: 130-140, 2015
- A.195 Nkizinkiko Y, Suneel Kumar BV, Jeankumar VU, Haikarainen T, Koivunen J, Madhuri C, Yogeewari P, Venkannagari H, Obaji E, Pihlajaniemi T, Sriram D, and Lehtiö L. Discovery of potent and selective nonplanar tankyrase inhibiting nicotinamide mimics. *Bioorg Med Chem* 23: 4139-4149, 2015
- A.196 Richter K, Konzack A, Pihlajaniemi T, Heljasvaara R, and Kietzmann T. Redox-fibrosis: Impact of TGFβ1 on ROS generators, mediators and functional consequences. *Redox Biol* 28: 344-352, 2015
- A.197 Tu H, Huhtala P, Lee H-M, Adams JC, and Pihlajaniemi T. Membrane-associated collagens with interrupted triple-helices (MACITs): evolution from a bilaterian common ancestor and functional conservation in *C. elegans*. *BMC Evol Biol* 15: 281, 2015
- A.198 Pirhonen J, Arola J, Sädevirta S, Luukkonen P, Karppinen SM, Pihlajaniemi T, Isomäki A, Hukkanen M, Yki-Järvinen H, and Ikonen E. Continuous grading of early fibrosis in NAFLD using label-free imaging: a proof-of-concept study. *PLoS One* 11: e0147804, 2016
- A.199 Karppinen SM, Honkanen H-K, Heljasvaara R, Riihilä P, Autio-Harmainen H, Sormunen R, Harjunen V, Väisänen M-R, Väisänen T, Hurskainen T, Tasanen-Määttä K, Kähäri V-

M, and Pihlajaniemi T. Collagens XV and XVIII show different expression and localization in cutaneous squamous cell carcinoma: type XV appears in tumour stroma while XVIII becomes up-regulated in tumour cells and lost from microvessels. *Exp Dermatol*, in press

Commentary: Lakshmanachetty S, and Koster MI. Emerging Roles for Collagen XV and XVIII in Cancer Progression. *Exp Dermatol*, in press

- A.200 Lin CHS, Chen J, Zhang Z, Johnson G, Cooper AJL, Feola J, Bank A, Shein J, Ruotsalainen H, Pihlajaniemi T, and Goligorsky MS. Endostatin and transglutaminase 2 are geronic proteins involved in fibrosis of aging kidney, *Kidney Internat*, in press
- A.201 Honkanen H-K, Izzi V, Holopainen T, Pihlajaniemi T, Alitalo K, and Heljasvaara R. Excess of VEGF-D modulates tumor inflammation and reduces growth of carcinogen-induced skin tumors. *Neoplasia*, 2016 (in revision)
- A.202 Kivinen N, Felszeghy S, Kinnunen AI, Setälä N, Aikio M, Kinnunen K, Sironen R, Pihlajaniemi T, Kauppinen A, and Kaarniranta K. Absence of collagen XVIII in mice causes age-related insufficiency in retinal pigment epithelium proteostasis. *Biogerontology* (in revision)
- A.203 Géraud C, Koch P-S, Damm F, Zierow J, Demory A, Dieltz M, Sticht C, Klapproth K, Busch K, Rodewald H-R, Breitkopf K, Richter K, Karppinen S-M, Pihlajaniemi T, Arnold B, Schledzewski K, and Goerdts S. Endothelial Gata4 controls organ-specific angiogenesis and liver development. *Submitted*
- A.204 Venkannagari H, Verheugd P, Koivunen J, Haikarainen T, Obaji E, Narwal M, Pihlajaniemi T, Lüscher B, and Lehtiö L. Chemical probe rescues cells from ARTD10 induced apoptosis and sensitizes cancer cells to DNA damage. *Submitted*
- A.205 Nathubhai A, Haikarainen T, Koivunen J, Hayward PC, Muñoz-Descalzo S, Tosh D, Lloyd MD, Koumanov F, Holman GD, Pihlajaniemi T, Lehtiö L, and Threadgill MD. Increasing insulin-activated glucose uptake, inhibiting Wnt signaling and in vitro anti-proliferative activity using extremely potent and selective tankyrase inhibitors. Manuscript in preparation.
- A.206 Izzi V, Kaur I, Devarajan R, Lakkala J, Savolainen E-R, Koistinen P, Ruotsalainen H, Prus P, Heljasvaara R, and Pihlajaniemi T. Collagen XVIII gatekeeps normal myeloid-erythroid balance and its deficiency characterizes acute myeloid leukemia. Manuscript in preparation.
- A.207 Heljasvaara R, Aikio M, and Pihlajaniemi T. Collagen XVIII in tissue homeostasis. *Matrix Biology*, Invited review in preparation.
- A.208 Martínez Nieto GA, Petäistö T, Liu H, Karppinen S-M, Ruotsalainen H, Kähäri V-M, Pihlajaniemi T, Donald Gullberg D, and Heljasvaara R. Role of integrin $\alpha 11$ in cutaneous squamous cell carcinoma. Manuscript in preparation.
- A.209 Vicente D, Izzi V, Finnilä M, Buler M, Mäkelä K, Aikio M, Karppinen SM, Herzig KH, Heljasvaara R, and Pihlajaniemi T. Specific isoforms of collagen XVIII affect hepatic glucose and lipid uptake and storage. Manuscript in preparation.
- A.210 Riihilä P, Nissinen L, Farshchian M, Kallajoki M, Kivisaari A, Meri S, Grénman R, Peltonen S, Peltonen J, Pihlajaniemi T, Heljasvaara R, and Kähäri V-M. Complement

Factor B and C3 regulate growth of cutaneous squamous cell carcinoma. Manuscript in preparation.

- A.211 Nkizinkiko Y, Desantis J, Koivunen J, Haikarainen T, Murthy S, Miele M, Obaji E, Pihlajaniemi T, Tabarrini O, and Lehtiö L. Tankyrase inhibitors as kinase inhibitors: double agents or off targets? Manuscript in preparation.
- A.212 Harjunen V*, Honkanen HK*, Youssef KK, Kinnunen A, Seppinen L, Boumahdi S, Aikio M, Brideau G, Lefebvre S, Mikkola ML, Glumoff V, Sormunen R, Blanpain C, Pihlajaniemi T, Heljasvaara R. Collagen XVIII is an essential extracellular component of the epidermal stem cell niche and regulates stem cell maintenance, hair follicle cycling and skin tumor formation.*Equal contribution. Manuscript in preparation.
- A.213 Härönen H, Zainul Z, Tu H, Shakirzyanova A, Sormunen R, Miinalainen I, Abdullin A, Naumenko N, Oikarainen T, Koski A, Martin P, Santoleri S, Koistinaho J, Silman I, Giniatullin R, Heikkinen A, and Pihlajaniemi T. Collagen XIII is indispensable for the neuromuscular junction organization and function, and shedding of its transmembrane form contributes to the synaptic plasticity. Manuscript in preparation.
- A.214 Kaur I, Ruskamo S, Koivunen J, Lackmann JJ, Petäjä-Repo U, Kursula P, Heljasvaara R, and Pihlajaniemi T. Biochemical and biophysical characterization of the aminoterminal domain of unknown function (DUF). Manuscript in preparation.

B Non-refereed scientific articles

Non-refereed journal article; book section; non-refereed conference proceedings

- B.1 Tryggvason K, and Pihlajaniemi T. Tyvikalvojen rakenne ja aineenvaihdunta (Structure and metabolism of basement membranes). *Duodecim* 96: 1532-1542, 1980
- B.2 Pihlajaniemi T. Perinnöllisten kollageenitautien molekyylibiologia (Molecular biology of inherited collagen disorders). *Duodecim* 104: 1802-1816, 1988
- B.3 Brideau G, Mäkinen M, Elamaa H, Tu H, Nilsson G, Alitalo K, Pihlajaniemi T, and Heljasvaara R. Endostatiini hillitsee imusuonten muodostusta ja imusolmukemetastaaseja hiirimallissa (Endostatin Overexpression Inhibits Lymphangiogenesis and Lymph Node Metastasis in Mice). *Duodecim* 124: 365, 2008
- B.4 Rasi K, Pihlajaniemi T, and Eklund L. Collagen XV deficiency predisposes to cardiomyopathy. Kollageenin XV puutos altistaa kardiomyopatiale. (Finnish Medical Society Journal, article in Finnish) *Duodecim* 126:2628, 2010
- B.5 Aikio M, Elamaa H, Vicente D, Izzi V, Kaur I, Seppinen L, Speedy HE, Kaminska D, Kuusisto S, Sormunen R, Heljasvaara R, Jones EL, Muilu M, Jauhiainen M, Pihlajaniemi T, Savolainen MJ, Shoulders CC, and Pihlajaniemi T. Tyvikalvokollageeni osallistuu rasvasolujen erilaistumiseen ja toimintaan. (Finnish Medical Society Journal, article in Finnish) *Duodecim* 130:1661, 2014.
- B.6 Pihlajaniemi T. Soluväliaineen tutkimuksen monet ulottuvuudet (Review: Many dimensions of extracellular matrix research). *Duodecim*, 129: 2262-2272, 2013

C Scientific books (monographs)

Book; edited book, conference proceedings or special issue of a journal

- C.1 Tryggvason K, Pihlajaniemi T, Salo T, Liotta LA, and Kivirikko KI. Studies on the biosynthesis and degradation of type IV procollagen. In: *New Trends in Basement Membrane Research*. Eds. Kuehn K, Schoene H, and Timpl R. Raven Press, New York, 187-193, 1982
- C.2 Tryggvason K, Pihlajaniemi T, and Salo T. Studies on the molecular composition and degradation of type IV procollagen. In: *Basement Membranes and Cell Movement*. Eds. Porter R, Whelan J. The Ciba Foundation, London, 117-124, 1984
- C.3 Prockop DJ, and Pihlajaniemi T. Collagen in normal and diseased connective tissue. In: *Arthritis and Allied Conditions*. Ed. McCarty DJ. Lea & Fabiger, Philadelphia, 10th edition, 210-226, 1984
- C.4 Prockop DJ, and Pihlajaniemi T. Collagen in normal and diseased connective tissue. In: *Arthritis and Allied Conditions*. Ed. McCarthy DJ. Lea & Fabiger, Philadelphia, 11th edition, 222-239, 1989
- C.5 Kivirikko KI, Myllylä R, and Pihlajaniemi T. Hydroxylation of proline and lysine residues in collagens and other animal and plant proteins. In: *Post-translational Modifications of Proteins*. Eds. Crabbe MJC, and Harding J. CRC Press, Boca Raton, 1-51, 1991
- C.6 Pihlajaniemi T. Type XIII collagen. In: *Human Protein Data*. Ed. Haeberli A. VCH Verlagsges. mbH, Weinheim, 2. Installment, 1993
- C.7 Pihlajaniemi T. Collagen type XV. In: *Human Protein Data*. Ed. Haeberli A. VCH Verlagsges. mbH, Weinheim, 3. Installment, 1995
- C.8 Pihlajaniemi T. Molecular properties of the glomerular basement membrane. In: *Molecular Pathology and Genetics of Alport Syndrome. Contributions to Nephrology*. Ed. Tryggvason K. Karger, Basel, 46-79, 1996
- C.9 Snellman A, and Pihlajaniemi T. Collagenous transmembrane proteins. In: *Recent research developments in biological chemistry*. Ed. Pandalai SG. Research Signpost, India, pp. 1 – 22, 2003
- C.10 Heljasvaara R, and Pihlajaniemi T. Experimental tumour models in mice. In: *Mouse as a Model Organism – from Animals to Cells*. Ed. Brakebusch C, and Pihlajaniemi T. Springer, pp. 89-104, 2011
- C.11 Brakebusch C, and Pihlajaniemi T. Foreword. In: *Mouse as a Model Organism – from Animals to Cells*. Ed. Pihlajaniemi T, and Brakebusch C. Springer, pp. 89-104, 2011

D Publications intended for professional communities

Article in a trade journal; article in a professional manual or guide or professional information system, textbook material; professional conference proceedings; published development or research report or study; textbook, professional manual or guide, dictionary

- D.1 Pihlajaniemi T. Väitöskirja akateemisena opinnäytteenä. In: *Kohti karonkkaa. Neuvoja väitöskirjaa tekeville*. Patja K, Huhtaniemi I, Ikonen E & Kontula K., eds, Kustannus Oy Duodecim, Helsinki, pp. 19-33, 1999
- D.2 Pihlajaniemi T. Innovative life sciences in Europe's north. *European Public Review*, 18: 208-209, 2009
- D.3 Pihlajaniemi T. Comments of the scientific director. Biocenter Oulu yearly reports, 1995–2008
- D.4 Pihlajaniemi T. Biotekniikan rahoituksen merkitys Suomen lääketieteen tutkimukselle. – *Duodecim* 123: 1371-1373, 2007

E Publications intended for the general public, linked to the applicant's research

Popularised article, newspaper article; popularised monograph

- E.1 Pihlajaniemi T. Elämäkertakirjoitus akatemiaprofessori Kari Kivirikosta. In: *Kansallisbiografia*, Helsinki. (In Swedish: Biografiskt lexikon; Kari Kivirikko Biografiskt lexikon för Finland), vol. III (Republiken), 2001
- E.2 Pihlajaniemi T. Kolumni-kirjoituksia (7 kappaletta). Sanomalehti Kaleva (Regular Columns for the Newspaper Kaleva), 2006-2007
- E.3 Pihlajaniemi T. Kari Kivirikko: Kollageenin synnyn salojen paljastaja. In: *Suomalaisia tieteen huipulla. Sata saavutusta*. Havaste P, Tiitta A & Markkanen, T., eds. Gaudeamus, pp. 166-167, 2014.

F Public artistic and design activities

Published independent work of art; public partial realisation of a work of art; public artistic performance or exhibition; model or design adopted for production/use

-

G Theses

- G.1 Pihlajaniemi T. Characterization of normal and diseased human collagens by the techniques of recombinant DNA. Doctoral dissertation, *Acta Universitatis Ouluensis*, A 189, 1987

H Patents and invention disclosures

- H.1 Synthesis of human procollagens and collagens in recombinant DNA systems (issued patent no: 5,593,859, U.S.A.; patent no: 2,121,698, Canada; patent no: AU 680,606, Australia; patent no: 99201209.6-2405, European patent; patent no: 0625048, Austria)
- H.2 Collagen-based methods and formulations for the treatment of arthritis, issued patent no: 5,925,736, U.S.A.;
- H.3 α 2 subunit of prolyl-4-hydroxylase, nucleic acid sequences encoding such subunit and methods for producing the same, patent no: 5,928,922, U.S.A.; patent no: 727281, Australia
- H.4 Method for diagnosing diseases, patent application no: 60/918,103, U.S.A.; OU10001-(collagen XV). Patent application no. 10176727.5, Europe.